



## NADAL<sup>®</sup> Strep A plus Test

Test cassette with star mould for the detection of group A streptococci

### Group A Streptococci

Beta-haemolytic group A streptococci (bacterial strain) are responsible for a range of different diseases. They are a major cause of infections of the upper respiratory tract, such as acute pharyngitis (tonsillitis) and scarlet fever. Streptococcal infections with streptococcus pyogenes are especially common amongst children aged 5-15 years. Due to their non-specific clinical symptoms, a distinction must be made between streptococcal and viral infections using appropriate diagnostics.

The early detection of streptococcal pharyngitis and the subsequent intervention is particularly important for children, as treatment can

- shorten the course of the disease
- ease the severity of symptoms
- reduce the likelihood of complications
- lower the chance of others being infected

### NADAL<sup>®</sup> Strep A Test

The NADAL<sup>®</sup> Strep A plus Test is an immunological rapid test for the fast, qualitative detection of group A streptococcus antigens in throat swabs, suitable for general practices, hospitals and laboratories. With a short testing time of just 5 minutes, as well as simple and precise results, the NADAL<sup>®</sup> Strep A plus Test is an ideal aid in the diagnosis of streptococcal infections.


### Benefits at a glance:

- ✓ High sensitivity and specificity
- ✓ Easy handling: results in 4 steps
- ✓ Reduction in use of unnecessary antibiotics
- ✓ Test storage at room temperature
- ✓ Minimum 12-month shelf life

Our NADAL<sup>®</sup> Strep A plus test incl. star mould:  
no more mixed up samples with this compact test  
system featuring an integrated reagent holder



## Test Procedure



Dispense **4 drops of reagent 1** into the extraction tube, followed by **4 drops of reagent 2**.

Place the **swab in the tube** and turn it against the sides of the tube using a circular motion. Repeat this at least 5 times.

Allow the solution to stand at room temperature for **1-2 minutes**. Press the swab firmly against the side of the tube and then remove it.

Using the dropper or disposable pipette, dispense **3 drops** of the extracted solution into the **sample well** of the test cassette.

Interpret test results after **5 minutes**.

A detailed description can be found in the package insert.

## Test Interpretation



## Strep A

Rapid test for the detection of group A streptococci in throat swabs

Product	Sample material	Sensitivity	Specificity	Product code	Pack size
NADAL® Strep A plus test cassette	Throat swab	97.6 %	97.5 %	222007	20 tests
NADAL® Strep A plus test cassette	Throat swab	97.6 %	97.5 %	222008	5 tests

Package includes: test cassettes • disposable pipette • extraction tube with dropper caps • sterile swab • extraction reagents 1 and 2 • positive controls

Rev01.01/XX0705en00033  
Rev01.01/EN05en00027

Our **customer service team** is at a hand to answer any questions you may have via telephone, email or in personal meetings. For details on **your local sales office and regional free phone numbers**, visit [www.nal-vonminden.com](http://www.nal-vonminden.com).

We help you care!

Rapid Tests

Laboratory Diagnostics

Laboratory Service

Consulting & Service

Headquarters

**nal von minden GmbH** · Carl-Zeiss-Strasse 12 · 47445 Moers · Germany

Chief Executive Officers: Sandra von Minden · Roland Meißner · Thomas Zander

Commercial reg. Kleve · HRB 5679

Tel.: +49 941 290 10-0 · Fax: +49 941 290 10-50

info@nal-vonminden.com  
[www.nal-vonminden.com](http://www.nal-vonminden.com)